

Demography of some non-native isopods (Crustacea, Isopoda, Oniscidea) in a Mid-Atlantic forest, USA

Elisabeth Hornung^{1,2}, Katalin Szlavecz², Miklós Dombos³

1 Department of Ecology, Institute for Biology, Faculty of Veterinary Science, Szent István University, H-1400 Budapest, P.O.Box 2, Hungary **2** Department of Earth and Planetary Sciences, The Johns Hopkins University, 3400 N. Charles St., Baltimore, MD 21218-2681, USA **3** Research Institute for Soil Science and Agricultural Chemistry of HAS, H-1525 Budapest, P.O. Box 102, Hungary

Corresponding author: *Elisabeth Hornung* (Hornung.Erzsebet@aotk.szie.hu)

Academic editor: *S. Taiti* | Received 16 February 2015 | Accepted 18 June 2015 | Published 30 July 2015

<http://zoobank.org/209651D5-D1FE-4161-9369-8BD5E888725A>

Citation: Hornung E, Szlavecz, K Dombos M (2015) Demography of some non-native isopods (Crustacea, Isopoda, Oniscidea) in a Mid-Atlantic forest, USA. In: Taiti S, Hornung E, Štrus J, Bouchon D (Eds) Trends in Terrestrial Isopod Biology. ZooKeys 515: 127–143. doi: 10.3897/zookeys.515.9403

Abstract

Introduced species dominate the terrestrial isopod fauna in most inland habitats of North America, including urban landscapes. These non-native species are often very abundant and thus potentially play a significant role in detritus processing. We monitored isopod assemblages in an urban forest for a year to examine the relationship between surface activity and abiotic environmental factors, and to analyze reproductive characteristics that might contribute to their successful establishment. Using pitfall trap samples we recorded five species, two of which, *Trachelipus rathkii* and *Cylisticus convexus*, were highly abundant. We determined size, sex and reproductive state of each individual. Surface activity of both species reflected variability in abiotic stress factors for isopods, such as soil moisture and soil temperature. Early spring the main trigger was soil temperature while later in the season increasing temperature and decreasing soil moisture jointly affected population dynamics. Activity significantly correlated with soil moisture. The temporal pattern of sex ratios supported the secondary sex ratio hypothesis. Males dominated the samples on the onset of the mating season in search of females. The pattern was reversed as females searched for suitable microsites for their offspring. Size independent fecundity decreased as conditions became more stressful late in the season.

Keywords

Abiotic drivers, activity density, reproductive patterns, secondary sex ratio hypothesis, urban soil fauna

Introduction

In recent years there has been an increased interest in non-native, expansive soil invertebrates in North America. Studies almost exclusively focused on earthworm invasion (e.g. James and Hendrix 2004, Hale et al. 2005, Szlavecz et al. 2011a). Earthworms, as ecosystem engineers have multiple, profound and visible effects on soil physical and biogeochemical processes. However, the soil decomposer food web is complex, and other members of the fauna also contribute to these processes. Soil invertebrate community structure with special emphasis of non-native species other than earthworms, has received less attention, partially because their effects on ecosystem processes may be more subtle (Niemelä et al. 1997).

Terrestrial isopods are macro-decomposers that can significantly contribute to detritus processing (comminution, inoculation) and nutrient release. They occur also in habitats too extreme for earthworms, such as salt marshes, arid grasslands and deserts. Here and in other habitats they can reach extremely high local densities (e.g. Paris and Pitelka 1962, Steinberger 1976, Sorensen and Burkett 1977, Shachak et al. 1979, Dias et al. 2005, Messina et al. 2012) elevating them to the rank of the primary detritivore grazers and keystone group in regulating fungal communities (Crowther et al. 2013).

About one-third of the North-American Oniscidea is non-native. The endemic species mostly concentrate in coastal areas, caves, and the southern regions of the continent (Leistikow and Wägele 1999, Jass and Klausmeier 2000). Introduction of non-native woodlice, mostly from Europe, has been going on for centuries. Many of these species are synanthropic, and, probably due to lack of native fauna, successfully invaded wildland habitats, agricultural fields and cities. Isopods are among the most abundant arthropods in urban landscapes (Bolger et al. 2000, Smith et al. 2006, Vilisics et al. 2007a). Undoubtedly, life history characteristics of successful species at least partially explain their dominant status.

In this paper we report data on isopod demography in an urban forest in Baltimore, Maryland, USA. The study was part of a larger ongoing monitoring effort coordinated by the Baltimore Ecosystem Study (www.beslter.org, BES thereafter). BES is one of the two urban sites within the Long Term Ecological Research (LTER) network in the USA. One overarching question BES explores how heterogeneity in social, physical and biological factors interact to influence biodiversity (including soil biodiversity) at multiple scales (Swan et al. 2011, Szlavecz et al. 2011b). In the present study we examined the relationship between isopod population characteristics and abiotic environmental factors, and further analyzed reproductive characteristics that might explain high local abundance and thus invasion success of the dominant species.

Methods

Study site

We surveyed the isopod fauna in Leakin Park, a 492 ha contiguous parkland in Baltimore, Maryland, USA (39°15'N, 76°30'W). The park is about 8 km NW from the urban core,

heavily forested, and surrounded with high density residential areas. The 90 year old forest belongs to the Tulip poplar Association (Brush et al. 1980) with common canopy species including Tulip poplar (*Liriodendron tulipifera*), several oaks (*Quercus alba*, *Q. coccinea*, *Q. velutina*), and American beech (*Fagus grandifolia*). Oaks and Tulip poplar make up 77% of the total annual litter mass, which is 4122 kg ha⁻¹ (Groffman et al. 2006). The soil belongs to the Legore series (fine-loamy, mixed, mesic, Ultic Hapludalf). Duff layer is thin (0–2 cm), bulk density and pH of A horizon are 1.12g cm⁻³, and 5.1, respectively. More detailed description of the vegetation and soils is given in Groffman et al. (2006).

The climate can be characterized by hot humid summers and cold winters with average annual air temperatures ranging from 14.5 °C in the inner urban areas to 12.8 °C in the surrounding rural areas. Precipitation is distributed evenly throughout the year in the region and ranges from an annual average of 106.8 cm in Baltimore to 103.1 cm in the surrounding metropolitan area (NOAA, www.nws.noaa.gov).

Sampling and laboratory measurements

Terrestrial isopods were sampled using pitfall traps (250 ml plastic cups) filled with propylene glycol. Ten traps were placed randomly around a 40 m × 40 m permanent forest plot established by the Baltimore Ecosystem Study LTER (Groffman et al. 2006). Traps operated between October 1999 and November 2000 and were emptied monthly except in late fall-winter when surface activity is generally low. The material was stored in 70% ethanol. All individuals were identified to species level using the nomenclature by Schmalfuss (2003).

Population and reproductive characteristics were determined only for the two abundant species. Because pitfall trap samples indicate a combination of surface activity and abundance of epigeic invertebrates, obtaining even relative density information creates a challenge. Recently, the term 'activity-density' has been used (Melbourne 1999, Westerman et al. 2008) to express abundance, and here we follow this practice. Activity density is expressed as number of individuals caught per trap per day.

We estimated body size by measuring the widest point of the head capsule (cephalon) at the level of the eyes (Sutton 1968). Measurements were taken to 0.01 mm accuracy under a dissecting stereo microscope. Adult females were divided into three reproductive categories: non-reproducing, gravid (either with eggs, embryos or mancas), and post-reproductive (with empty marsupium). Reproductive period was defined as the time span between the appearance of the first gravid females and that of the last one with brood pouch (marsupium) either with or without progeny (Sutton et al. 1984).

Abiotic factors

We obtained soil temperature and moisture data from the Baltimore Ecosystem Study database. Soil temperature was measured continuously using HOBO H8 Pro Series

Temp/External Temp data loggers at 10 cm depth. For soil moisture measurements six time domain reflectometry (TDR) waveguide probes (Soil Moisture Equipment Corporation) were installed vertically into the soil at random locations throughout the plot. The waveguide probes are 20 cm long, so those vertically installed span a depth of 0 to 20 cm below ground. Soil moisture was measured once every four to six weeks.

Statistical analysis

Mean daily values were used to explore correlation between soil temperature and activity density. Distance-weighted least squares fitting was used for smoothing. Due to lack of continuous soil moisture data, relationships between activity density and soil moisture was explored by using Spearman Rank Order Correlation Coefficient. We used multiple linear regression to explore relationships among fecundity (number of eggs produced by females), body size (cephalon width) and sampling date for each species. For computing these analyses we used STATISTICA 12 software (StatSoft Inc. 1984–2013).

To compare relative importance of independent variables i.e. size and time, we used standardized beta partial regression coefficients. Beta coefficients are obtained by setting all the variables to a mean of 0 and standard deviation to 1.

Ratio of males is expressed as the total number of males (M) over total number of adults (N) caught during a given trapping period (Dangerfield and Telford 1994). We conducted two different statistical tests. First, we wanted to estimate the unknown sex ratio and its uncertainty. Second, we wanted to assess the probability of the null hypothesis that the observed data is consistent with a M:N ratio of 0.5. A simple chi-square test informs only of the latter, moreover, chi-square test assumes Gaussian distribution i.e. symmetric error bars. This approximation is only valid when N is large. Since several of our samples were small, we chose a different approach (Gotelli and Ellison 2004). At a given sex ratio the distribution of observed M males out of a sample of N individuals ($M < N$) follows a binomial distribution. We first calculated the cumulative probability (CDF) that the number of males in the sample is equal or less than the measured count M. If M is more than half the sample size we estimate the probability that the number of males is equal or more than the count. Using this cumulative binomial distribution, we first computed the 95% confidence intervals in both directions, which often are not symmetric (Table 1). Finally we determined the probability the observed number of males is consistent with a sex ratio of 0.5, by simply reading off the midpoint value of the CDF.

Results

During the course of sixteen months a total of 2480 isopods were caught. The following five species were recorded: *Haplophthalmus danicus* Budde-Lund, 1880, *Hyloniscus riparius* (C. Koch, 1838), *Philoscia muscorum* (Scopoli, 1763), *Trachelipus rathkii* (Brandt, 1833) and *Cylisticus convexus* (De Geer, 1778). The pitfall material was domi-

Table 1. Abundance of *Trachelipus rathkii* and *Cylisticus convexus* between October 1999 and November 2000 in Leakin Park urban forest, Baltimore, USA. Numbers from all pitfall traps are pooled for each sampling period. Male ratio was calculated as proportion of males in total sample. 95% confidence intervals for these estimates are given the parentheses. Bold letters indicate significant differences from the expected 0.5 value.

Species	Month	Males	Females	Ratio of males	<i>p</i>
<i>Trachelipus rathkii</i>	October	110	118	0.48 (0.43; 0.54)	0.321
	November	38	61	0.38 (0.31; 0.47)	0.013
	January	2	2	0.50 (0.25; 0.91)	0.312
	March	9	25	0.26 (0.17; 0.42)	0.004
	April	76	58	0.57 (0.50; 0.64)	0.050
	May	462	445	0.51 (0.49; 0.54)	0.275
	June	244	306	0.44 (0.41; 0.48)	0.005
	July	401	461	0.47 (0.44; 0.50)	0.022
	August	141	230	0.38 (0.35; 0.43)	< 0.001
	September	90	128	0.41 (0.37; 0.48)	0.006
	October	16	21	0.43 (0.32; 0.58)	0.256
	November	2	4	0.33 (0.15; 0.73)	0.343
<i>Cylisticus convexus</i>	October	185	252	0.42 (0.39; 0.47)	0.001
	November	38	55	0.41 (0.34; 0.50)	0.048
	January	0	0	NA	NA
	March	4	8	0.33 (0.18; 0.62)	0.194
	April	11	20	0.35 (0.24; 0.52)	0.075
	May	178	315	0.36 (0.33; 0.40)	< 0.001
	June	263	353	0.51 (0.48; 0.55)	0.298
	July	352	568	0.38 (0.36; 0.41)	< 0.001
	August	353	163	0.61 (0.57; 0.65)	< 0.001
	September	162	227	0.42 (0.38; 0.46)	0.001
	October	48	38	0.56 (0.47; 0.65)	0.117
	November	6	16	0.27 (0.16; 0.47)	0.026

nated by the latter two species, with 1270 *T. rathkii* and 1073 *C. convexus* (53 and 45 %) individuals, respectively. Detailed analysis of the population characteristics is given only for these two species.

Surface activity, demographic changes

Isopod numbers began to increase late April, peaked in July and declined in September and ceased by November (Fig. 1A and B). Activity density of both species positively correlated with soil temperature ($r = 0.75$, and $r = 0.87$, for *T. rathkii*, and for *C. con-*




Figure 1. Temporal changes in activity-density of *Cylisticus convexus* (A) and *Trachelipus rathkii* (B) and soil physical characteristics in an urban forest in Baltimore. Mean numbers of individuals captured daily per trap (*C. convexus*: open circles, *T. rathkii*: asterix) \pm SE are shown. Dotted line: mean daily soil temperature at 10 cm; dashed line: smoothed soil temperature; open triangles: volumetric soil moisture content.

vexus, respectively). Soil moisture decreased during the growing season (Fig. 1A, B). Excluding winter months, when no isopods were caught, isopod activity positively correlated with soil moisture during the growing season (Spearman $R = 0.32$, $p < 0.05$ for *T. rathkii*, and $R = 0.35$, $p < 0.05$ for *C. convexus*).

Reproductive characters

Male ratio

Male ratio varied over time, with the highest and lowest male : total ratio being 0.57 and 0.26 for *T. rathkii*, 0.61 and 0.27 for *C. convexus*, respectively (Table 1). For *T. rathkii*, we detected significant deviations from the expected the 0.5 ratio in seven sampling dates. In all but two months (April, May) activity density of females exceeded that of males. For *C. convexus* we detected significant differences also in seven cases. Again, only two months did males exceed females, but this happened later in the season (August, October).

Reproductive period, phenology and fecundity

Trachelipus rathkii started reproducing late April – early May (Table 2) with 20% of the females being gravid, and all in the egg carrying stage. Proportion of gravid females peaked at 57% in June–mid-July and slightly declined (47%) by mid-August. Gravid *C. convexus* females appeared in the traps later in the season (June–July), but at this period females both with eggs and empty marsupium were present. In September only a single gravid *T. rathkii* was caught, while 16% of the 146 female *C. convexus* in the sample were still gravid, all in post-reproductive stage.

Fecundity of females in *C. convexus* and *T. rathkii* were compared over time. We analyzed the relationship between fecundity and body size using multiple linear regression models: clutch size (number of eggs), as dependent variable, and body size (head width) and time (days from the start of the investigation), as independent variables. Regression summary for *C. convexus*: adjusted $R^2 = 0.65$, $F(2,119) = 112.10$, $p < 0.001$; $\text{beta}(\text{day}) = -0.44$; $\text{beta}(\text{head width}) = 0.37$. Regression summary for *T. rathkii*: adjusted $R^2 = 0.38$, $F(2,123) = 38.836$, $p < 0.001$; $\text{beta}(\text{day}) = -0.22$; $\text{beta}(\text{head width}) = 0.65$. Number of eggs increased with body size (Figure 2A), and these relationships were sta-

Table 2. Percentage of reproductive *Trachelipus rathkii* and *Cylisticus convexus* in Leakin Park, Baltimore. Two stages are distinguished. Gravid: with eggs, embryos or mancas in the marsupium; postreproductive: empty marsupium.

			May	June	July	Aug
<i>Trachelipus rathkii</i>	All reproductive*		20.6	57.1	47.1	0.6
		Gravid**	100	41.7	17.5	0
		Postreproductive**	0	58.3	82.5	100
<i>Cylisticus convexus</i>	All reproductive*		0	61.0	40.0	16.0
		Gravid**	0	29.4	57.1	0
		Postreproductive**	0	70.6	42.9	100

*Calculated as percentage of all females in the sample

** Calculated as percentage of all reproductive females in the sample




Figure 2. Fecundity of the dominant isopod species in Leakin Park, Baltimore. **A–B:** Relationship between fecundity and size (**A**) and its stability over time (**B**) **C–D:** Change of fecundity over time (**C**) and size independent fecundity over time, based on the residuals of egg numbers (**D**). *Cylisticus convexus*: open circles and dashed lines; *Trachelipus nathkii*: crosses and dotted lines.

ble throughout the season (Figure 2B). Size independent fecundity decreased with time for both species (Figure 2C): residuals of egg numbers showed negative trends, however, their slopes were different (Figure 2D). Standardized beta partial regression coefficients were -0.44 ($p < 0.001$) and -0.22 for *C. convexus* and *T. rathkii*, respectively, indicating that the decrease of female fecundity during the season was not linked to body size.

Discussion

Species composition

In the Greater Baltimore Metropolitan Area we have recorded a total of eleven terrestrial isopod species (Hornung and Szlavetz 2003). All species are non-native, and most have been known from North America for a hundred years (Hatch 1947, van Name 1936, 1940, 1942, Eberly 1953, Lindroth 1957, Jass and Klausmeyer 2000). Urban fauna is often characterized by synanthropic, generalist species that may lead to higher community similarity among cities (McKinney 2006). All five species present in Leakin Park fit this category, and the two dominant species *Trachelipus rathkii* and *Cylisticus convexus*, are among the ten most abundant exotics in North America (Jass and Klausmeier 2000). Both species are characterized as expansive in Europe and have been introduced to other continents, as well (Gruner 1966). In Europe both species occur in a variety of habitats, including cities (Korsós et al. 2002; Hornung et al. 2007, 2008, Vilisics and Hornung 2009, Vilisics et al. 2007a,b). The difference between their occurrences in the two continents is that *C. convexus* tends to be more synanthropic in Central and Western Europe, its abundance is moderate to low, and is scarce or missing in the North (Gruner 1966, Berg et al. 2008, Vilisics et al. 2007a). In North America both species are widely distributed, and abundant populations of *C. convexus* and *T. rathkii* were reported e.g. from Michigan (Hatchett 1947) and Wisconsin (Jass and Klausmeier 1996), too.

Temporal patterns

Temperature and relative humidity are known to be the main drivers of terrestrial isopod activity (Warburg 1987, 1993). Our data support this statement, but also show a more complex relationship in the field. Soil temperature is the main factor triggering surface activity early spring; however, later in the season increasing temperature and decreasing soil moisture jointly affect dynamics of the populations. In early fall even though temperature remains high, activity decreases (Fig. 1). At this time soil moisture is low due to high evapotranspiration rates by trees and lack of precipitation (Groffman et al. 2006). We acknowledge that other factors, such as size, mobility, and behavior may bias pitfall trap samples. However, the huge number of individuals in the samples gives us confidence that the dynamics we detected for these two abundant species, is real.

Reproductive characteristics

Male ratio

With the exception of parthenogenetic species, where males occur in extremely low numbers, the sex ratio of most isopod species can be described by bimodality (e.g. Paris and Pitelka 1962, Sorensen and Burkett 1977, Hornung 1989, 1991). There are some examples for species with very different and constant adult sex ratio, e.g. *Porcellio ficulneus* Budde-Lund, 1885 male: female = 1:9 (Warburg 2007), and *Schizidium tiberianum* Verhoeff, 1923, 1:6 (Warburg and Cohen 1991). Sex ratio for terrestrial woodlice is routinely reported in papers focusing on population characteristics. In most cases these values are only snapshots obtained from samples of a short time period and do not reflect temporal changes of the sex ratio. Sex ratio might differ seasonally depending on different mortality and/or activity of sexes. We are aware of only a few long term studies where populations were frequently sampled to obtain changes of sex ratio over time (e.g. Paris and Pitelka 1962, Sorensen and Burkett 1977). It is therefore important to exercise caution when comparing data for different populations even for the same species, as variations both in space and time may occur. Montesanto et al. (2008) reported a special case for *Platyarthrus aiasensis* Legrand, 1954 comparing 19 populations in and around Sicily. The populations differed in male ratio from 0 (parthenogenetic) to 0.37 of males.

Deviation from the expected 0.5:0.5 ratio may be due to behavioral differences between the sexes especially during reproductive period as proposed by the secondary sex ratio hypothesis (Dangerfield and Hassall 1994). According to this hypothesis at the onset of the mating season males are more active looking for receptive females. Later, gravid females exceed males in the sample, as they are looking for favorable microhabitats that maintain optimal conditions for their progeny. Both *T. rathkii* and *C. convexus* follow this pattern in our study site; the difference between them is the peak of timing of male searching behavior. High male dominance (0.98) was found also for *Protracheoniscus politus* (C. Koch, 1841) in Hungary (Oberfrank et al. 2011) and for *Armadillidium vulgare* (Latreille, 1804) in Texas (Sorensen and Burkett 1977) at the beginning of the activity season, before the onset of reproduction.

Gruner (1966) compiled and qualitatively reported population data for the two species we studied here. The percentage of *C. convexus* males was found to be lower than 50% in Denmark, France and Italy. For *T. rathkii* a female predominance was also reported.

Reproductive period and phenology

In Europe the onset of the reproductive period varied with latitude for both *C. convexus* and *T. rathkii* (Gruner 1966). For instance, *C. convexus* started reproducing in April in France; in Denmark the onset shifted to June. The number of offspring ranged between

14–50 per female, with extremely high numbers (73 embryos per female) in Italy (Gruner 1966). *T. rathkii* was reported to have two broods between May and September.

Jass and Klausmeier (2004) found a latitudinal difference in reproductive peaks in North America for several species including the ones studied here. In Wisconsin gravid females of *C. convexus* were present during June–July in the populations while reproductive period of *T. rathkii* was longer (June–August). In the present study the reproductive period started earlier, in April both for *T. rathkii*, and for *C. convexus*, and ended in August and September, respectively.

Reproductive output over time

Reproductive output is an important component of life history strategies and has a cost of decreased parental survival (Stearns 1976). The strong correlation between female size and egg number in terrestrial isopods is well known (e.g. Dangerfield and Telford 1990, Ma et al. 1991, Nair 1978, Warburg 2011). However, females may not allocate the same amount of energy into reproduction during the entire season. We found that the mean number of eggs/embryos decreased independently of female size as the reproductive period progressed. This tendency is the consequence of increasing environmental stress (unfavorable humidity and temperature changes, drought) and contributes to the costs of survival (Cody 1966, Charnov 2002, Hornung and Warburg 1993, 1994, 1998). Isopods can either invest less or regain part of the energy invested in reproduction earlier by reabsorbing ovarian oocytes or marsupial eggs under stressful conditions (Hornung and Warburg 1993, 1994).

Successful establishment, expansion, invasion

There is still some confusion regarding terminology in invasion ecology. The term invasive species is used for species “exhibiting rapid spread, irrespective to impact” (Davis 2009), as well as for species showing “demonstrable ecological or economic impact” (Lockwood et al. 2007). The two isopod species we reported here undoubtedly fit the first description, as both are well established and common throughout North America (Jass and Klausmeier 2000). The main characteristics successful invaders share include high dispersal rate, high genetic variability, short generation time, large number of offspring, broad diet and ecological tolerance. Some of these traits overlap with what Sutton et al. (1984) describes for “eurodynamic”, essentially *r* strategist species. Detritivores are by definition resource generalists (even though they have food preferences) allowing them to find food in a variety of habitats. Both *C. convexus* and *T. rathkii* have high fecundity, and can reproduce several times per season. Moreover, female isopods have been shown to be able to store sperm and utilize their stock in repeated reproduction events (Suzuki and Ziegler 2005). This can be especially significant when only one or a few females are transported from one habitat to another. Isopods are relatively easy

to transport with soil, mulch, ornamental plants, timber and other means. Once introduced, they can quickly establish and, being mobile epigeic species, further disperse on their own. Garthwaite et al. (1995) estimated that *Armadillidium vulgare*, another common species in North America spread across the continent and reached the West Coast in about 150 years. We can only speculate that the lack of native relatives or ecologically equivalent soil fauna further facilitated the spread of terrestrial isopods including the species studied here.

The highly altered urban habitats can serve both as points of introduction via trade or transportation, and refuges for non-indigenous soil fauna. Residential areas provide food (e.g. compost, mulch) and shelter (building foundation, landscaping objects), while green corridors or even underground pipe systems can be conduits for dispersal. High epigeic isopod abundance has been repeatedly shown in urban habitat fragments and suburbs (Bolger et al. 2000, Smith et al. 2006, Vilisics et al. 2007a). Often, the same species dominate urban isopod fauna. *T. rathkii* and *C. convexus* belong to this successful cosmopolitan group, contributing to the global phenomenon of biotic homogenization in cities (Perrings et al. 2010).

Conclusions

We examined population dynamics and reproductive characteristics of terrestrial isopods in an urban forest in Baltimore, Maryland, USA. Temporal patterns of male ratio supported the secondary sex ratio hypothesis for both dominant species. As expected, fecundity was correlated with female size. However, size independent reproductive output declined during the active season indicating a response to increasing stress. High fecundity, good dispersal ability and broad habitat and resource tolerance all may contribute to the invasion success of the investigated species in North America. Additionally, lack of native competitors and locally favorable conditions can further facilitate their spread and persistence in many ecosystems.

Acknowledgements

This study was supported by grants from the National Science Foundation (DEB 9714835), the Hungarian Science Foundation (MTA 049-OTKA 31623 and OTKA T043508 to EH and KS). We thank undergraduate students at Johns Hopkins University for their help in the field and in the lab. Dan Dillon compiled the temperature and moisture data, and Alex Szalay helped with the statistics. EH gratefully acknowledges the hospitality of the Dept. of Earth and Planetary Sciences during her stay in the USA.

EH processed the samples in the laboratory, identified and measured the isopods, and determined their reproductive state. MD carried out the statistical tests. KS did all field work and sorted the pitfall trap samples. All three authors contributed to writing the manuscript.

We thank two anonymous reviewers for their helpful comments on an earlier version of the manuscript. This paper was supported by the 9877-3/2015/FEKUT grant of the Hungarian Ministry of Human Resources.

References

- Berg MP, Soesbergen M, Tempelman D, Wijnhoven H (2008) Verspreidingsatlas Nederland landpissebedden, duizendpoten miljoenpoten (Isopoda, Chilopoda, Diplopoda). EIS-Nederland, Leiden & Vrije Universiteit – Afdeling Dierecologie, Amsterdam, 192 pp.
- Bolger DT, Suarez AV, Crooks KR, Morrison SA, Case TJ (2000) Arthropods in urban habitat fragments in southern California: area, age, and edge effects. *Ecological Applications* 10: 1230–1248. doi: 10.1890/1051-0761(2000)010[1230:AIUHFI]2.0.CO;2
- Brush GS, Lenk C, Smith J (1980) The natural forests of Maryland: an explanation of the vegetation map of Maryland. *Ecological Monographs* 50: 77–92. doi: 10.2307/2937247
- Cody ML (1966) A general theory of clutch size. *Evolution* 20: 174–184. doi: 10.2307/2406571
- Charnov EL (2002) Reproductive effort, offspring size and benefit-cost ratios in the classification of life histories. *Evolutionary Ecology Research* 4: 749–758. <http://hdl.handle.net/1928/1653>
- Crowther TW, Stanton WG, Thomas SM, A’Bear AD, Hiscox J, Jones H, Voříšková S, Baldrian P, Boddy L (2013) Top-down control of soil fungal community composition by a globally distributed keystone consumer. *Ecology* 94(11): 2518–2528. doi: 10.1890/13-0197.1
- Dangerfield JM, Hassall M (1994) Shelter use and secondary sex ratios in the woodlice *Armadillidium vulgare* and *Porcellio scaber* (Crustacea: Isopoda). *Journal of Zoological Society of London* 233: 1–7. doi: 10.1111/j.1469-7998.1994.tb05257.x
- Dangerfield JM, Telford SR (1990) Breeding phenology, variation in reproductive effort and offspring size in a tropical population of the woodlouse *Porcellionides pruinosus*. *Oecologia* 82: 251–258. doi: 10.1007/BF00323542
- Dangerfield JM, Telford SR (1994) Population size structure and sex ratios in some woodlice (Crustacea: Oniscidae) from Southern Africa. *Journal of Tropical Ecology* 10(2): 261–271. doi: 10.1017/S0266467400007902
- Davis MA (2009) *Invasion biology*. Oxford University Press, Oxford, 244 pp.
- Dias N, Sprung M, Hassall M (2005) The abundance and life histories of terrestrial isopods in a salt marsh of the Ria Formosa lagoon system, southern Portugal. *Marine Biology* 147: 1343–1352. doi: 10.1007/s00227-005-0033-2
- Eberly W (1953) The terrestrial isopods of Indiana. *Proceedings of the Indiana Academy of Science* 63: 272–277.
- Garthwaite RL, Lawson R, Sassaman C (1995) Population genetics of *Armadillidium vulgare* in Europe and North America. In: Alikhan MA (Ed.) *Terrestrial Isopod Biology*. Proceedings of the International Symposium of Terrestrial Isopod Biology. Crustacean Issues 9. A.A. Balkema, Rotterdam, 145–199.
- Gotelli NJ, Ellison AM (2004) *A Primer of Ecological Statistics*. Sinauer Associates, Inc., Sunderland MA, 510 pp.

- Groffman P, Pouyat RV, Cadenasso ML, Zipperer WC, Szlavecz K, Yesilonis IC, Band LE, Brush GS (2006) Land use context and natural soil controls on plant community composition and soil nitrogen and carbon dynamics in urban and rural forests. *Forest Ecology and Management* 236: 177–192. doi: 10.1016/j.foreco.2006.09.002
- Gruner HE (1966) Krebstiere oder Crustacea. 5. Isopoda. In: Dahl M, Preuss F (Eds) *Die Tierwelt Deutschlands und der angrenzenden Meeresteile und ihre Lebensweise*. Gustav Fischer Verlag, Jena, 156–158.
- Hale CM, Frelich LE, Reich PB (2005) Exotic european earthworm invasion dynamics in northern hardwood forests of Minnesota, USA. *Ecological Applications* 15(3): 848–860. doi: 10.1890/03-5345
- Hatch M (1947) The Chelifera and Isopoda of Washington and adjacent regions. University of Washington, Publications in Biology 10: 155–274. [Oniscidea pp. 174–205]
- Hatchett SP (1947) Biology of the Isopoda of Michigan. *Ecological Monographs* 17: 48–79. doi: 10.2307/1948613
- Hornung E (1989) Population dynamics and spatial distribution of *Trachelipus nodulosus* (Koch CL, 1838) (Crustacea Isopoda) in a sandy grassland. In: Ferrara FAR, Manicacstri C, Schmalfluss H, Taiti S (Eds) *Monitore Zoologico Italiano (N.S.)*. Il Sedicesimo, Firenze. Monographs 4, 399–409.
- Hornung E (1991) Isopod distribution in a heterogeneous grassland habitat. In: Juchault P, Mocquard JP (Eds) *Third Symposium on the Biology of Terrestrial Isopods*. Universiteit de Poitiers, France, 7–79.
- Hornung E (2011) Evolutionary adaptation of oniscidean isopods to terrestrial life: Structural – physiological – behavioural aspects. *Terrestrial Arthropod Reviews* 4(2): 95–130. doi: 10.1163/187498311X576262
- Hornung E, Szlavecz K (2003) Establishment of a Mediterranean isopod (*Chaetophiloscia sicula* Verhoeff, 1908) in a North American temperate forest. *Crustaceana Monographs* 2: 181–189. <http://www.brill.com/biology-terrestrial-isopods>
- Hornung E, Tóthmérész B, Magura T, Vilisics F (2007) Changes of isopod assemblages along an urban-suburban-rural gradient in Hungary. *European Journal of Soil Biology* 43: 158–161. doi: 10.1016/j.ejsobi.2007.01.001
- Hornung E, Vilisics F, Sólymos P (2008) Low alpha and high beta diversity in terrestrial isopod assemblages in the Transdanubian region of Hungary. In: Zimmer M, Cheikrouha C, Taiti S (Eds) *Proceedings of the International Symposium of Terrestrial Isopod Biology - ISTIB-7*. Shaker Verlag, Aachen, 1–13.
- Hornung E, Warburg MR (1993) Breeding patterns in the oniscid isopod, *Porcellio ficulneus* Verh., at high temperature and under different photophases. *Invertebrate Reproduction and Development* 23: 151–158. doi: 10.1080/07924259.1993.9672306
- Hornung E, Warburg MR (1994) Oosorption and oocyte loss in *Porcellio ficulneus* B.-L. (Isopoda; Oniscidea; Porcellionidae) under stressful conditions. *Tissue & Cell* 26: 277–284. doi: 10.1016/0040-8166(94)90102-3
- Hornung E, Warburg MR (1998) Plasticity of a *Porcellio ficulneus* population under extreme weather conditions (a case study). *Israel Journal of Zoology* 44: 395–398.

- James SW, Hendrix PF (2004) Invasion of exotic earthworms into North America and other regions. In: Edwards CA (Ed.) *Earthworm Ecology*. CRC Press, Boca Raton, 75–88.
- Jass J, Klausmeier B (1996) Comparison of Wisconsin terrestrial isopods and their life cycle traits. *Field Station Bulletin Milwaukee, Wisconsin* Fall 29(2): 14–26.
- Jass J, Klausmeier B (2000) Endemics and immigrants: North American terrestrial isopods (Isopoda, Oniscidea) North of Mexico. *Crustaceana* 73(7): 771–799. doi: 10.1163/156854000504804
- Jass J, Klausmeier B (2004) The terrestrial isopod *Hyloniscus riparius* (Isopoda: Oniscidea: Trichoniscidae) in Wisconsin. *The Great Lake Entomologist* 36(1–2): 70–75.
- Korsós Z, Hornung E, Szilávecz K, Kontschán J (2002) Isopoda and Diplopoda of urban habitats: New data to the fauna of Budapest. *Annales Historico-Naturales Musei Nationalis Hungarici* 94: 45–51.
- Leistikow A, Wägele JW (1999) Checklist of the terrestrial isopods of the new world Crustacea, Isopoda, Oniscidea). *Revista brasileira de Zoologia* 16(1): 1–72. doi: 10.1590/S0101-81751999000100001
- Lindroth CH (1957) *The faunal connections between Europe and North America*. John Wiley & Sons, Inc., New York, 344 pp. doi: 10.5962/bhl.title.6759
- Lockwood JL, Hoopes MF, Marchetti MP (2013) *Invasion ecology*. John Wiley & Sons, 456 pp.
- Ma HHT, Lam PKS, Dudgeon D (1991) Inter- and intraspecific variation in the life histories of three sympatric isopods in a Hong Kong forest. *Journal of Zoology, London* 224: 677–687. doi: 10.1111/j.1469-7998.1991.tb03795.x
- McKinney ML (2006) Urbanisation as major cause of biotic homogenization. *Biological Conservation* 127: 247–260. doi: 10.1016/j.biocon.2005.09.005
- Melbourne B (1999) Bias in the effect of habitat structure on pitfall traps: An experimental evaluation. *Australian Journal of Ecology* 24(3): 228–239. doi: 10.1046/j.1442-9993.1999.00967.x
- Messina G, Pezzino E, Montesanto G, Caruso G, Lombardo BM (2012) The diversity of terrestrial isopods in the natural reserve “Saline di Trapani e Paceco” (Crustacea, Isopoda, Oniscidea) in northwestern Sicily. *ZooKeys* 176: 215–230. doi: 10.3897/zookeys.176.2367
- Montesanto G, Caruso D, Lombardo BM (2008) Genetic variability in parthenogenetic and amphigonic populations of *Platyarthrus aiasensis* Legrand from Sicily (Isopoda: Oniscidea). In: Zimmer M, Charfi-Cheikhrouha F, Taiti S (Eds) *Proceedings of the International Symposium of Terrestrial Isopod Biology – ISTIB 7*. Shaker Verlag, Aachen.
- Nair GA (1978) Some aspects of the population characteristics of the soil Isopod *Porcellio laevis* (Latreille), in Delhi region. *Zoologische Anzeiger, Jena* 201(1–2): 86–96.
- Niemelä J, Spence JE, Cárcamo H (1997) Establishment and interactions of carabid populations: an experiment with native and introduced species. *Ecography* 20: 643–652. doi: 10.1111/j.1600-0587.1997.tb00433.x
- NOAA (2015) National Oceanic and Atmospheric Administration. http://www.nws.noaa.gov/climate/local_data.php?wfo=lwx [verified on 10/06/2015]
- Oberfrank A, Végh A, Lang Z, Hornung E (2011) Reproductive strategy of *Protracheoniscus politus* (C.Koch, 1841) (Oniscidea: Crinocheta: Agnaridae). In: Zidar P, Štrus J (Eds) Pro-

- ceedings of the 8th International Symposium of Terrestrial Isopod Biology, Bled, Slovenia, 135–136.
- Paris OH, Pitelka FA (1962) Population characteristics of the terrestrial isopod *Armadillidium vulgare* in California grassland. *Ecology* 43: 229–248. doi: 10.2307/1931979
- Perrings C, Mooney H, Williamson M (2010) *Bioinvasion & Globalization. Ecology, Economics, Management, and Policy.* Oxford University Press Inc., New York, 267 pp.
- Schmalfuss H (2003) World catalog of terrestrial isopods (Isopoda: Oniscidea). *Stuttgarter Beiträge zur Naturkunde, Serie A* 654: 1–341.
- Shachak M, Steinberger Y, Orr Y (1979) Phenology, activity and regulation of radiation load in the desert isopod, *Hemilepistus reaumuri*. *Oecologia* 40(2): 133–140. doi: 10.1007/BF00347931
- Smith RM, Warren PH, Thompson K, Gaston KJ (2006) Urban domestic gardens (VI): environmental correlates of invertebrate species richness. *Biodiversity and Conservation* 15: 2415–2438. doi: 10.1016/j.biocon.2005.10.045
- Sorensen EMB, Burkett RD (1977) A population study of the isopod, *Armadillidium vulgare*, in Northeastern Texas. *The Southwestern Naturalist* 22(3): 375–387. <http://www.jstor.org/stable/30054805>
- Stearns SC (1976) Life-history tactics: a review of the ideas. *Quarterly Review of Biology* 51(1): 3–47. doi: 10.1086/409052
- Steinberger Y (1976) Feeding, energy flow and soil turnover in the desert isopod, *Hemilepistus reaumuri*. *Oecologia* 24: 57–69. doi: 10.1007/BF00545487
- StatSoft, Inc. (2003) STATISTICA (data analysis software system), version 6. www.statsoft.com
- Sutton SL (1968) The population dynamics of *Trichoniscus pusillus* and *Philoscia muscorum* (Crustacea, Oniscidea) in limestone grassland. *Journal of Animal Ecology* 37(2): 425–444. doi: 10.2307/2958
- Sutton SL, Hassall M, Willows R, Davis RC, Grundy A, Sunderland KD (1984) Life histories of terrestrial isopods: a study of intra- and interspecific variation. *Symposia of the Zoological Society of London* 53: 269–294.
- Suzuki S, Ziegler A (2005) Structural investigation of the female genitalia and sperm-storage sites in the terrestrial isopod *Armadillidium vulgare* (Crustacea, Isopoda). *Arthropod Structure & Development* 34: 441–454. doi:10.1016/j.asd.2005.06.002
- Swan CM, Pickett STA, Szlavecz K, Warren PS, Willey KT (2011) Biodiversity and community composition in urban ecosystems: coupled human, spatial and metacommunity processes. In: Niemelä J, Breuste JH, Guntenspergen G, McIntyre NE, Elmqvist T, James P (Eds) *Urban Ecology: Patterns, Processes, and Applications.* Oxford University Press, 179–186. doi: 10.1093/acprof:oso/9780199563562.003.0021
- Szlavecz K, McCormick M, Xia L, Saunders J, Morcol T, Whigham D, Filley T, Csuzdi Cs, (2011a) Ecosystem effects of non-native earthworms in Mid-Atlantic deciduous forests. *Biological Invasions* 15: 1165–1182. doi: 10.1007/s10530-011-9959-0
- Szlavecz K, Warren P, Pickett S (2011b) Biodiversity on the Urban Landscape, Chapter 6. In: Cincotta RP, Gorenflo LJ (Eds) *Human Population: Its Influences on Biological Diversity.* Ecological Studies 214: 78–101. doi: 10.1007/978-3-642-16707-2_6

- van Name W (1936) The American land and freshwater isopod Crustacea. *Bulletin of the American Museum of natural History* 71: 1–535.
- van Name W (1940) A supplement to the American land and freshwater isopod Crustacea. *Bulletin of the American Museum of natural History* 77: 109–142.
- van Name W (1942) A second supplement to the American land and freshwater isopod Crustacea. *Bulletin of the American Museum of natural History* 80: 299–329.
- Vilisics F, Elek Z, Lovei GL, Hornung E (2007a) Composition of terrestrial isopod assemblages along an urbanisation gradient in Denmark. *Pedobiologia* 51: 45–53. doi: 10.1016/j.pedobi.2006.12.004
- Vilisics F, Sólymos P, Hornung E (2007b) A preliminary study on habitat features and associated terrestrial isopod species. In: Tajovský K, Schlaghamerský J, Pižl V (Eds) *Contributions to Soil Zoology in Central Europe II*, 195–199.
- Vilisics F, Hornung E (2009) Urban areas as hot-spots for introduced and shelters for native isopod species. *Urban Ecosystems* 12: 333–345. doi: 10.1007/s11252-009-0097-8
- Warburg MR (1987) Isopods and their terrestrial environment. *Advances in Ecological Research* 22: 161–172. doi:10.1016/S0065-2504(08)60246-9
- Warburg MR (1993) *Evolutionary biology of land isopods*. Springer Verlag, Berlin-Heidelberg, 159 pp. doi: 10.1007/978-3-662-21889-1
- Warburg MR (2007) Distribution, reproduction, and relative abundance of Oniscids: a long-term study on isopods (Isopoda, Oniscidea) in Israel. *Crustaceana* 80(10): 1223–1252. doi: 10.1163/156854007782321218
- Warburg MR (2011) Cost of breeding in Oniscid Isopods: a partial review. *Crustaceana* 84(12–13): 1561–1580. doi: 10.1163/156854011X607006
- Warburg MR, Cohen N (1991) Reproductive pattern, allocation, and potential in a semelparous isopod from the Mediterranean region of Israel. *Journal of Crustacean Biology* 11(3): 368–374. doi: 10.2307/1548463
- Westerman PR, Borza JK, Andjelkovic J, Liebman M, Danielson B (2008) Density-dependent predation of weed seeds in maize fields. *Journal of Applied Ecology* 45: 1612–1620. doi: 10.1111/j.1365-2664.2008.01481.x